

BALLS IN THE TRIANGULAR RATIO METRIC

S. HOKUNI, R. KLÉN, Y. LI, AND M. VUORINEN

ABSTRACT. We consider the triangular ratio metric and estimate the radius of convexity for balls in some special domains and inclusion relations of metric balls defined by the triangular ratio metric, the quasihyperbolic metric and the j -metric.

1. INTRODUCTION

Geometric function theory makes use of several metrics for subdomains of $\mathbb{R}^n$. It has turned out that sometimes hyperbolic metric or more generally, hyperbolic type metrics, see [Kl3, L], are more natural than the Euclidean metric, because of their better invariance or quasi-invariance properties. On the other hand, hyperbolic type metrics are sometimes difficult to estimate and it is desirable to find simple expressions to serve as comparison functions. Two such expressions, the angular metric and the triangular ratio metric, were recently studied in [KLW]. Here our goal is to study the triangular ratio metric further for proper subdomains of $\mathbb{R}^n$. In particular, we study the local convexity of balls in this metric for some simple domains. For some other metrics, results of this type were recently proven by R. Klén in [Kl2, Kl4], in answer to a question posed in [Vu2].

We study also inclusion relations between balls in different metrics. We are interested the triangular ratio metric, the quasihyperbolic metric and the j -metric. We consider the inclusion relations in general domains as well as in some specific examples as the punctured space the the half-space. These kind of results for hyperbolic type metrics have been studied in [KV, KV2].

For a domain $G \subsetneq \mathbb{R}^n$, and $x, y \in G$, we define the triangular ratio metric by

$$(1.1) \quad s_G(x, y) = \sup_{z \in \partial G} \frac{|x - y|}{|z - x| + |z - y|} \in [0, 1],$$

the j -metric by

$$j_G(x, y) = \log \left(1 + \frac{|x - y|}{\min\{d_G(x), d_G(y)\}} \right),$$

2010 *Mathematics Subject Classification.* Primary 51M10; Secondary 52A20 .

Key words and phrases. Local convexity, metric ball, triangular ratio metric, quasihyperbolic metric, j -metric.

File: hklv121210b.tex, printed: 2012-12-12, 2.24

where $d_G(z) = d(z, \partial G)$, and the quasihyperbolic metric by

$$k_G(x, y) = \inf_{\gamma} \int_{\gamma} \frac{|dz|}{d_G(z)},$$

where the infimum is taken over all rectifiable curves $\gamma \subset G$ joining x and y . G.J. Martin and B.G. Osgood proved in [MO, page 38] the following formula for the quasihyperbolic distance:

$$k_G(x, y) = \sqrt{\alpha^2 + \log^2 \frac{|x|}{|y|}}, \quad x, y \in G = \mathbb{R}^n \setminus \{0\},$$

where $\alpha = \angle(x, 0, y)$. For a metric space (G, m) we define the metric ball for $x \in G$ and $r > 0$ by $B_m(x, r) = \{y \in G : m(x, y) < r\}$.

In Sections 3-5 we consider local convexity properties of balls in triangular ratio metric. In Section 3 we consider the punctured space $\mathbb{R}^n \setminus \{0\}$, in Section 4 the half-space $\mathbb{H}^n$ and in Section 5 the punctured half-space $\mathbb{H}^n \setminus \{e_n\}$ and polygons $P \subset \mathbb{R}^2$. In Sections 6 and 7 we study the inclusion of balls defined by the triangular ratio metric, the quasihyperbolic metric and the j -metric. In Section 6 we consider these metrics in general domains and in Section 7 in the punctured space and in the half-space.

Our main results are the following two theorems.

Theorem 1.2. 1) Let $G = \mathbb{R}^n \setminus \{z\}$, $z \in \mathbb{R}^n$, $x \in G$ and $r > 0$. The metric ball $B_s(x, r)$ is (strictly) convex if $r \leq 1/2$ ($r < 1/2$).
2) Let $x \in G = \mathbb{H}^n$ and $r \in (0, 1)$. Then

$$B_s(x, r) = B^n \left(x - e_n x_n \left(1 - \frac{1+r^2}{1-r^2} \right), \frac{2x_n r}{1-r^2} \right)$$

and $B_s(x, r)$ is thus strictly convex.

3) Let $x = (x_1, x_2) \in G = \mathbb{H}^2 \setminus \{e_n\}$ with $x_2 < |x_1|$ and $r \in (0, r_0]$, where

$$r_0 = \frac{\sqrt{x_1^2 + x_2^2} - \sqrt{2}x_2}{|x_1| + x_2}.$$

Then $B_s(x, r)$ is convex.

4) Let $P \subset \mathbb{R}^2$ be a polygon and $x \in P$. Then $B_s(x, r)$ is convex for all $r \in (0, 1/2]$. Moreover, if P is convex then $B_s(x, r)$ is convex for all $r \in (0, 1)$.

Theorem 1.3. 1) Suppose that $G \subset \mathbb{R}^n$ is a domain. For each $x \in G$ and $r \in (0, 1)$, we have

$$B^n \left(x, \frac{2r}{1+r} d_G(x) \right) \subset B_s(x, r) \subset B^n \left(x, \frac{2r}{1-r} d_G(x) \right).$$

2) Suppose that $G \subset \mathbb{R}^n$ is a domain. For each $x \in G$ and $r \in (0, 1)$, we have

$$B_j(x, \log(1 + 2r)) \subset B_s(x, r) \subset B_j\left(x, \log\left(1 + \frac{2r}{1 - 3r}\right)\right)$$

and the first inclusion is sharp if there exists some point w in $\partial B_s(x, r)$ such that $d_G(x) = d_G(w)$ and

$$\partial G \cap S^{n-1}(w, d_G(w)) \cap S^{n-1}(x, d_G(x)) \neq \emptyset.$$

3) Let $G = \mathbb{R}^n \setminus \{0\}$. For each $x \in G$ and $r \in (0, 1)$, we have

$$B\left(x, \frac{2r}{1+r}|x|\right) \subset B_s(x, r) \subset B\left(x, \frac{2r}{1-r}|x|\right).$$

Moreover, the inclusions are sharp.

4) Let $G = \mathbb{R}^n \setminus \{0\}$. For each $x \in G$ and $r \in (0, 1)$, we have

$$B_j(x, m) \subset B_s(x, r) \subset B_j(x, M),$$

where $m = \log(1 + 2r)$, and $M = \log(1 + \frac{2r}{1-r})$. Moreover, the inclusions are sharp.

2. PRELIMINARY RESULTS

Fact 2.1. Let us have a polar function $r = r(\theta)$. Then the slope of the curve $r = r(\theta)$ at the point (r, θ) is

$$\frac{r(\theta) + \tan(\theta) \frac{dr}{d\theta}}{-r(\theta) \tan(\theta) + \frac{dr}{d\theta}}.$$

By the definition of s (1.1) the supremum is attained at a point z that is on a maximal ellipsoid with foci at x and y . Therefore it is clear that s is monotone with respect to the domain, which means that if $G \subset \mathbb{R}^n$ and $G' \subset G$ are domains and $x, y \in G'$ then $s_G(x, y) \leq s_{G'}(x, y)$.

By the definition and the monotonicity with respect to domain it is easy to prove that for all $x \in G$ and $r \in (0, 1)$

$$(2.2) \quad B_{s_G}(x, r) = \bigcap_{z \in \partial G} B_{s_{\mathbb{R}^n \setminus \{z\}}}(x, r).$$

We denote the hyperbolic distance by ρ_G , where G is either the unit ball $\mathbb{B}^n$ or the upper half space $\mathbb{H}^n$.

Lemma 2.3. Let $a, b > 0$.

(1) The function $f_1(r) \equiv \frac{\log(1+ar)}{r}$ is strictly decreasing from $(0, \infty)$ onto $(0, a)$.

(2) The function $f_2(r) \equiv \frac{r}{2-r} - \operatorname{ar} \tanh r$ is strictly decreasing from $(0, 1)$ onto $(-\infty, 0)$.

(3) The function $f_3(r) \equiv \frac{r}{r(r-2)\log(1-r)}$ is strictly decreasing from $(0, 1)$ onto $(0, 1/2)$.

(4) The function $f_4(t) \equiv \log t - \frac{t-1}{t+1}$ is increasing from $(1, \infty)$ onto $(0, \infty)$.

Proof. (1) Let $f_{11}(r) = \log(1 + ar)$ and $f_{12}(r) = r$. It is clear that $f_{11}(0^+) = f_{12}(0^+) = 0$. By differentiation,

$$\frac{f'_{11}(r)}{f'_{12}(r)} = \frac{a}{1 + ar}$$

which is strictly decreasing in $(0, \infty)$. Therefore, we get the monotonicity of f_1 . The limiting values are clear by l'Hôpital's rule.

(2) By differentiation,

$$f'_2(r) = \frac{-3r^2 + 4r - 2}{(1 - r^2)(2 - r)^2}$$

which is negative. Therefore, f_2 is strictly decreasing. The limiting values are clear.

(3) Let $f_{31}(r) = \frac{r}{2-r}$ and $f_{32}(r) = -\log(1 - r)$. It is clear that $f_{31}(0^+) = f_{32}(0^+) = 0$. By differentiation,

$$\frac{f'_{31}(r)}{f'_{32}(r)} = \frac{2(1 - r)}{(2 - r)^2} \equiv 2\phi(r).$$

By differentiation, we have

$$\phi'(r) = -\frac{r}{(2 - r)^3} < 0.$$

Then ϕ is strictly decreasing in $(0, 1)$. Therefore, we get the monotonicity of f_3 . The limiting values are clear by l'Hôpital's rule.

(4) By differentiation,

$$f'_4(t) = \frac{t(1 + t)^2}{t^2 + 1} > 0,$$

Therefore, f_4 is strictly increasing. The limiting values are clear. $\square$

3. CONVEXITY OF BALLS IN PUNCTURED SPACE

We consider the metric s in the punctured space $\mathbb{R}^n \setminus \{0\}$. We first compare s with the hyperbolic metric ρ and then study convexity of metric balls $B_s(x, r)$.

By the definition it is clear that for $x, y \in G = \mathbb{R}^n \setminus \{0\}$, we have

$$s_G(x, y) = \frac{|x - y|}{|x| + |y|}.$$

This special case of the triangular ratio metric has been studied in [B, AST].

Theorem 3.1. *The inequality $s_G(x, y) \leq \frac{1}{\log 3} j_G(x, y)$ holds for all $x, y \in G = \mathbb{R}^n \setminus \{0\}$.*

Proof. Without loss of generality, we may assume that $|x| \leq |y|$. Then for all $x, y \in G$, by Lemma 2.3 (1), we have

$$\frac{j_G(x, y)}{s_G(x, y)} = \frac{\log \left(1 + \frac{|x-y|}{|x|} \right)}{\frac{|x-y|}{|x|+|y|}} \geq \log \left(1 + \frac{|x|+|y|}{|x|} \right) \geq \log 3.$$

Equality holds if $0, x, y$ are collinear and $|x| = |y|$. $\square$

Open problem 3.2. We notice that $s_G(x, y) \leq \frac{1}{\log 3} j_G(x, y) < j_G(x, y)$ for $G = \mathbb{R}^n \setminus \{0\}$. A natural problem is that is it true that $s_G \leq j_G$ holds for all subdomain $G \subset \mathbb{R}^n$?

We only get the following constant which is larger than 1 but less than 2.

Theorem 3.3. Suppose that $G \subset \mathbb{R}^n$ is a domain. Then for all $x, y \in G$ we have

$$s_G(x, y) \leq \frac{1}{\log 2} j_G(x, y).$$

Proof. For given $x, y \in G$, we may assume that $d_G(x) \leq d_G(y)$. Then if $|x - y| > d_G(x)$, obviously we have

$$s_G(x, y) \leq 1 \leq \frac{1}{\log 2} j_G(x, y).$$

If $|x - y| \leq d_G(x)$, then

$$s_G(x, y) \leq \frac{|x - y|}{2d_G(x)} \leq \log \left(1 + \frac{|x - y|}{d_G(x)} \right) = j_G(x, y).$$

$\square$

Let us next consider convexity of metric balls.

Lemma 3.4. For all $\alpha \in (0, \arccos(1 - 2r^2))$ and $r \in (0, 1/2]$ we have

$$1 + \cos \alpha - 2\sqrt{2} \cos(\alpha/2) \sqrt{\cos \alpha + 2r^2 - 1} \geq 0.$$

Proof. Let us write previous inequality in a following form

$$g(\alpha) = \cos \alpha - 2\sqrt{2} \cos(\alpha/2) \sqrt{\cos \alpha + 2r^2 - 1} \geq -1.$$

First we need to show that $g(\alpha)$ is strictly increasing. For this we need

$$(3.5) \quad g'(\alpha) = \sin \alpha \left(\frac{\sqrt{2}(r^2 + \cos \alpha)}{\cos(\alpha/2) \sqrt{\cos \alpha + 2r^2 - 1}} - 1 \right).$$

In order to show that $g'(\alpha) > 0$ we need to prove following inequality

$$h(\alpha) = \frac{\sqrt{2}(r^2 + \cos \alpha)}{\cos(\alpha/2) \sqrt{\cos \alpha + 2r^2 - 1}} > 1.$$

Because

$$h'(\alpha) = \frac{\sqrt{2} \tan(\alpha/2) (r^2 - 1)^2}{\cos(\alpha/2) (\cos \alpha + 2r^2 - 1)^{3/2}}$$

it is easy to see that $h'(\alpha) > 0$ for all $\alpha \in (0, \arccos(1 - 2r^2))$ and $r \in (0, 1/2]$. Since $h(\alpha)$ is strictly increasing we see that $h(0) < h(\alpha)$ and with a simple computation we get

$$h(\alpha) > h(0) = r + 1/r \geq \frac{5}{2} > 1.$$

Now we have shown that $g(\alpha)$ is strictly increasing. Therefore $g(\alpha) \geq g(0) = 4r - 1 > -1$ and the assertion follows. $\square$

Theorem 3.6. *Let $G = \mathbb{R}^n \setminus \{z\}$, $z \in \mathbb{R}^n$, $x \in G$ and $r \in (0, 1]$. The metric ball $B_s(x, r)$ is nonconvex if $r > 1/2$.*

Proof. By symmetry it is sufficient to consider only the case $n = 2$, $z = 0$ and $x = 2$. Let $r \in (0, 1]$, $y \in \partial B_s(x, r)$ and denote $t = |y|$. Now $s(x, y) = r$ is equivalent to

$$|2 - y| = (2 + t)r.$$

By the law of cosines we obtain $|2 - y|^2 = t^2 + 4 - 4t \cos \alpha$ which is equivalent to

$$|2 - y| = \sqrt{t^2 + 4 - 4t \cos \alpha}.$$




FIGURE 1. Parametrization of $\partial B_s(x, r)$ in the proof of Theorem 3.6.

Therefore we get

$$\sqrt{t^2 + 4 - 4t \cos \alpha} = (2 + t)r$$

and by solving t with respect to α we obtain

$$(3.7) \quad t^*(\alpha) = \frac{2 \left(r^2 + \cos \alpha \pm \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right)}{1 - r^2}.$$

In this proof we select

$$t(\alpha) = \frac{2 \left(r^2 + \cos \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right)}{1 - r^2}.$$

Next we solve the following inequality to check, which values of α are interesting

$$(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1) > 0.$$

With a simple substitution $a = \cos \alpha$ we get

$$a^2 + 2r^2t + 2r^2 - 1 > 0 \Leftrightarrow a > 1 - 2r^2 \Leftrightarrow \alpha < \arccos(1 - 2r^2).$$

It is clear that if $r > 1/2$, then $\arccos(1 - 2r^2) > \pi/3$. Therefore it is enough to focus on angles $\alpha \in (0, \pi/3)$. The slope of the tangent of $\partial B_s(x, r)$ with respect to α according to the Definition 2.1 is

$$\begin{aligned} m(\alpha) &= \frac{t(\alpha) + \tan \alpha \, t'(\alpha)}{-t(\alpha) \tan \alpha + t'(\alpha)} \\ &= \frac{\sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} + \sin \alpha \tan \alpha}{\sin \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \tan \alpha}, \end{aligned}$$

where

$$t'(\alpha) = \frac{-2 \left(r^2 + \cos \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right) \sin \alpha}{(r^2 - 1) \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)}}.$$

Since $B_s(2, r)$ is symmetric with respect to x_1 -axis, we need to show that $m(\alpha) < 0$ for some $\alpha \in (0, \pi/3)$. It is clear that for all $\alpha \in (0, \pi/3)$ inequality

$$\sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} + \sin \alpha \tan \alpha > 0$$

holds. Now we need to show that for some $\alpha \in (0, \pi/3)$ inequality

$$f(\alpha) = \sin \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \tan \alpha < 0$$

holds. With a simple computation we see that $f(0) = 0$ and $f'(0) = 1 - 2r < 0$ for all $r > \frac{1}{2}$. Therefore $f(\alpha) < 0$ for some sufficiently small α . Now we have shown that $m(\alpha) < 0$ for some $\alpha \in (0, \pi/3)$ and the assertion follows. $\square$

Theorem 3.8. *Let $G = \mathbb{R}^n \setminus \{z\}$, $z \in \mathbb{R}^n$, $x \in G$ and $r > 0$. The metric ball $B_s(x, r)$ is (strictly) convex if $r \leq 1/2$ ($r < 1/2$).*

Proof. By symmetry it is sufficient to consider only the case $n = 2$, $z = 0$ and $x = 2$. Let us use following markings obtained in (3.7)

$$\begin{aligned} t_1(\alpha) &= \frac{2 \left(r^2 + \cos \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right)}{1 - r^2} \\ t_2(\alpha) &= \frac{2 \left(r^2 + \cos \alpha + \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right)}{1 - r^2} \end{aligned}$$

In the proof of Theorem 3.6 we showed that for $t_1(\alpha)$ the slope of the tangent of $\partial B_s(x, r)$ is

$$m_1(\alpha) = \frac{\sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} + \sin \alpha \tan \alpha}{\sin \alpha - \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \tan \alpha}$$

and $m_1(0) < 0$ if $r > 1/2$. With this it is easy to see that

$$\begin{aligned} m_1(0) &> 0 \quad \text{if } r < 1/2, \\ m_1(0) &\rightarrow \infty \quad \text{if } r \rightarrow 1/2. \end{aligned}$$

Similarly, for $t_2(\alpha)$ we write

$$t_2'(\alpha) = \frac{2 \left(r^2 + \cos \alpha + \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \right) \sin \alpha}{(r^2 - 1) \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)}}$$

and

$$\begin{aligned} m_2(\alpha) &= \frac{t_2(\alpha) + \tan \alpha \, t_2'(\alpha)}{-t_2(\alpha) \tan \alpha + t_2'(\alpha)} \\ &= \frac{-\sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} + \sin \alpha \tan \alpha}{\sin \alpha + \sqrt{(1 + \cos \alpha)(\cos \alpha + 2r^2 - 1)} \tan \alpha}, \end{aligned}$$

where $m_2(\alpha) \rightarrow -\infty$ if $\alpha \rightarrow 0$. To prove the theorem we need to show that $m_1'(\alpha) \leq 0$ and $m_2'(\alpha) \geq 0$ for all $\alpha \in (0, \arccos(1 - 2r^2))$ where $r \in (0, 1/2]$. Firstly, we get

$$m_1'(\alpha) = \frac{-2r^2 \cos^2(\alpha/2) \frac{1}{\cos^2 \alpha} \left(\varphi(\alpha) - 2\sqrt{2} \sqrt{\cos^2(\alpha/2) \omega(\alpha)} \right)}{\sqrt{\varphi(\alpha) \omega(\alpha)} \left(\sin \alpha - \tan \alpha \sqrt{\varphi(\alpha) \omega(\alpha)} \right)^2},$$

where $\varphi(\alpha) = 1 + \cos \alpha$ and $\omega(\alpha) = \cos \alpha + 2r^2 - 1$. Because

$$\varphi(\alpha) \omega(\alpha) > 0 \Leftrightarrow \alpha < \arccos(1 - 2r^2),$$

it is easy to see that for all $\alpha \in (0, \arccos(1 - 2r^2))$ and $r \in (0, 1/2]$ the following inequality

$$\sqrt{\varphi(\alpha) \omega(\alpha)} \left(\sin \alpha - \tan \alpha \sqrt{\varphi(\alpha) \omega(\alpha)} \right)^2 > 0$$

holds. To prove that $m_1'(\alpha) \leq 0$ we need to show that the inequality

$$-2r^2 \cos^2(\alpha/2) \frac{1}{\cos^2 \alpha} \left(\varphi(\alpha) - 2\sqrt{2} \sqrt{\cos^2(\alpha/2) \omega(\alpha)} \right) \leq 0$$

holds. It is easy to see that

$$-2r^2 \cos^2(\alpha/2) \frac{1}{\cos^2 \alpha} < 0$$

for all $\alpha \in (0, \arccos(1 - 2r^2))$ and $r \in (0, 1/2]$. In Lemma 3.4 we showed that

$$g(\alpha) = 1 + \cos \alpha - 2\sqrt{2} \sqrt{\cos^2(\alpha/2) (\cos \alpha + 2r^2 - 1)} \geq 0.$$

Therefore $m'_1(\alpha) \leq 0$. For $m_2(\alpha)$ we get

$$m'_2(\alpha) = \frac{2r^2 \cos^2(\alpha/2) \frac{1}{\cos^2 \alpha} \left(\varphi(\alpha) + \sqrt{2} \sqrt{\cos(2\alpha) + 4r^2 \cos \alpha + 4r^2 - 1} \right)}{\sqrt{\varphi(\alpha)\omega(\alpha)} \left(\sin \alpha + \tan \alpha \sqrt{\varphi(\alpha)\omega(\alpha)} \right)^2},$$

Because

$$\cos(2\alpha) + 4r^2 \cos \alpha + 4r^2 - 1 \geq 0 \Leftrightarrow \alpha < \arccos(1 - 2r^2),$$

we can clearly see that $m'_2(\alpha) > 0$ for all $\alpha \in (0, \arccos(1 - 2r^2))$ and the assertion follows. $\square$




FIGURE 2. Contours of the metric balls $B_s(x, r)$ in $\mathbb{R}^2 \setminus \{0\}$ where $x = 2$ and $r = 0.2, 0.4, 0.5$ and 0.65 .


4. CONVEXITY OF BALLS IN HALF-SPACE

We consider the triangular ratio metric in half space $\mathbb{H}^n$. We compare first s with ρ and then prove that the metric balls $B_s(x, r)$ are also Euclidean and thus always convex.

By the definition we obtain that for $x, y \in G = \mathbb{H}^n$

$$(4.1) \quad s_G(x, y) = \frac{|x - y|}{|x - y'|},$$

where $y' = (y_1, \dots, y_{n-1}, -y_n) \notin G$, see Figure 3.

FIGURE 3. Formula for s in $\mathbb{H}^n$ as in (4.1).

Theorem 4.2. *The inequality $s_{\mathbb{H}^n}(x, y) \leq \rho_{\mathbb{H}^n}(x, y)$ holds for all $x, y \in \mathbb{H}^n$.*

Proof. It suffices to consider the 2-dimension case. We assume that $\text{Im } x \leq \text{Im } y$. Let $z = [x, \bar{y}] \cap \partial\mathbb{H}^2$, $\alpha = \angle(z, x, y)$ and $\beta = \angle(z, y, x)$. It is easy to see that $0 \leq \beta \leq \alpha \leq \pi$. Then

$$s_{\mathbb{H}^2}(x, y) = \frac{|x - y|}{|x - z| + |z - y|} = \frac{\sin(\alpha + \beta)}{\sin \alpha + \sin \beta} = \frac{t - 1}{t + 1}$$

and

$$\rho_{\mathbb{H}^2}(x, y) = \log |x_*, x, y, y_*| = \log t,$$

where $t = \frac{\cos \frac{\alpha}{2} \cos \frac{\beta}{2}}{\sin \frac{\alpha}{2} \sin \frac{\beta}{2}} \in (1, \infty)$. By Lemma 2.3 (4), we have

$$s_{\mathbb{H}^2}(x, y) \leq \rho_{\mathbb{H}^2}(x, y).$$

This completes the proof. □

Lemma 4.3. *Let $z, y_1 \in \mathbb{R}$, $y_2, a > 0$ and*

$$f(z) = \sqrt{z^2 + a^2} + \sqrt{(y_1 - z)^2 + y_2^2}.$$

Then $f\left(\frac{ay_1}{y_2 + a}\right) \leq f(z)$ for all $z \in \mathbb{R}$.

Proof. With a simple computation we obtain

$$f'(z) = \frac{z - y_1}{\sqrt{y_2^2 + (y_1 - z)^2}} + \frac{z}{\sqrt{a^2 + z^2}}$$

and $f'(z) = 0$ is equivalent to $z = \frac{ay_1}{y_2 + a}$. To prove this lemma we focus on three different cases.

Case 1. Let us assume that $y_1 > 0$. Now $\hat{z} > 0$ and $0 < \hat{z} < y_1$. With a simple computation we get $f'(0) = -\frac{y_1}{\sqrt{y_1^2 + y_2^2}} < 0$ and $f'(y_1) = \frac{y_1}{\sqrt{y_1^2 + a^2}} > 0$.

Case 2. Let us assume that $y_1 < 0$. Now $\hat{z} < 0$ and $y_1 < \hat{z} < 0$. With a simple computation we get $f'(0) = -\frac{y_1}{\sqrt{y_1^2 + y_2^2}} > 0$ and $f'(y_1) = \frac{y_1}{\sqrt{y_1^2 + a^2}} < 0$.

Case 3. Finally, let us assume that $y_1 = 0$. Now $\hat{z} = 0$ and $-1 < \hat{z} < 1$. With a simple computation we get $f'(1) = \frac{1}{\sqrt{1+a^2}} + \frac{1}{\sqrt{y_2^2+1}} > 0$ and $f'(-1) = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{y_2^2+1}} < 0$. $\square$

Theorem 4.4. *Let $x \in G = \mathbb{H}^n$ and $r \in (0, 1)$. Then*

$$B_s(x, r) = B^n \left(x - e_n x_n \left(1 - \frac{1+r^2}{1-r^2} \right), \frac{2x_n r}{1-r^2} \right).$$

Proof. By symmetry of the domain it suffices to consider only the case $n = 2$ and we may assume that $x = (0, a)$, $a > 0$. First we select points $q = (0, q_2)$ and $w = (0, w_2)$ such that $s(x, q) = s(x, w) = r$ where $q_2 \in (0, a)$, $w_2 > a$ and $r \in (0, 1)$. By the definition of s we get for q_2

$$\frac{a - q_2}{a + q_2} = r$$

which is equivalent for $q_2 = \frac{a-ar}{1+r}$. In a similar way we obtain $w_2 = \frac{a-ar}{-1-r}$. With a simple computation we get

$$|C| = \frac{w_2 + q_2}{2} = \frac{a(1+r^2)}{1-r^2}$$

$$R = \frac{w_2 - q_2}{2} = \frac{2ar}{1-r^2}.$$

Next we choose $y = (y_1, y_2)$ such that $y_1 \in \mathbb{R}$ and $y_2 > 0$. To prove




FIGURE 4. Illustration of the Theorem 4.4.

the theorem we need to show that $s(x, y) = r$ if and only if $|C - y| = R$ for $C = \frac{a(1+r^2)}{1-r^2}i$. By using polar coordinates to y with respect to the

center point C and radius R we obtain

$$y = C + Re^{it} = \left(\frac{2ar}{1-r^2} \cos t, \frac{a(1+r^2)}{1-r^2} + \frac{2ar}{1-r^2} \sin t \right), \quad t \in [0, 2\pi).$$

Next we need to find the point $z \in \partial G$ which gives the supremum in the definition of s . In other words we need to maximize following function

$$f(z) = \frac{|x - y|}{\sqrt{z^2 + a^2} + \sqrt{(y_1 - z)^2 + y_2^2}}.$$

By using Lemma 4.3 we obtain that $z = \frac{ay_1}{y_2 + a}$ maximizes function $f(z)$. By transforming previously obtained polar coordinates to z we get

$$z = \frac{aR \cos t}{a + |C| + R \sin t} = \frac{ar \cos t}{1 + r \sin t}.$$

By using these polar coordinate expressions of y and z we can write

$$s(x, y) = f(z) = \frac{\frac{2ar}{(1-r^2)} \sqrt{1+r^2+2r \sin t}}{\frac{a \sqrt{(1+r^2+2r \sin t)^3}}{(1-r^2)(1+r \sin t)} \frac{a \sqrt{1+r^2+2r \sin t}}{1+r \sin t}} = r$$

and the assertion follows. $\square$




FIGURE 5. Contours of the metric balls $B_s(x, r)$ in $\mathbb{H}^2$ where $x = (0, 1)$ and $r = 0.2, 0.4, 0.5$ and 0.6 .

Corollary 4.5. *Let $x \in G = \mathbb{H}^n$ and $r \in (0, x_n)$. Then*

$$B^n(x, r) = B_s \left(x - e_n \left(x_n - \sqrt{x_n^2 - r^2} \right), \frac{x_n - \sqrt{x_n^2 - r^2}}{r} \right).$$

Corollary 4.6. *Let $x \in G = \mathbb{H}^n$ and $r > 0$. Then*

$$B^n \left(x, \frac{\sqrt{r^2 + x_n^2} - x_n}{r} \right) = B_s \left(x - e_n \left(x_n - \sqrt{x_n^2 + r^2} \right), r \right).$$

Corollary 4.7. *Let $x \in \mathbb{H}^n$ and $r \in (0, 1)$. Then*

$$B_s(x, r) = B_\rho(x - e_n x_n (1 - a), t), \\ t = \operatorname{ar tanh} \left(\frac{2r}{1+r^2} \right), \quad b = \frac{(1+r^2)}{\cosh(t)(1-r^2)}.$$

Proof. By symmetry of the domain it suffices to prove the result in the case $n = 2$ for $x = (0, a)$. By [Vu1, 2.2] and Theorem 4.4

$$B_\rho(ai, t) = B^2(a \cosh(t)i, a \sinh(t)) = B^2 \left(\frac{a(1+r^2)}{1-r^2}i, \frac{2ar}{1-r^2} \right),$$

which is equivalent to

$$\begin{cases} a \cosh t = \frac{x_2(1+r^2)}{1-r^2} \\ a \sinh t = \frac{2x_2 r}{1-r^2} \end{cases}$$

and the assertion follows. $\square$

5. CONVEXITY OF BALLS IN PUNCTURED HALF-SPACE AND POLYGONS

We consider the triangular ratio metric in the punctured half space $G = \mathbb{H}^n \setminus \{e_n\}$. By (2.2), Theorems 3.8 and 4.4 it is clear that the following result holds.

Lemma 5.1. *Let $x \in G = \mathbb{H}^n \setminus \{e_n\}$ and $r \in (0, 1/2]$. Then $B_s(x, r)$ is convex.*

However, the upper bound for the radius r in Lemma 5.1 is not sharp. To see this we can choose x close to $\partial\mathbb{H}^n$ and far from e_n . Now $B_s(x, r)$ is a Euclidean ball even for $r \in (1/2, r_0]$. For example for $x = e_n/10$ it can be verified that $B_s(x, r)$ is convex for $r \in (0, 3/4]$, see Figure 6.

The disks $B_s(x, r)$ for $x \in \mathbb{R}^2 \setminus \{e_2\}$ and large enough radius consists of two parts separated by curve $\{(t, (t^2 + 1)/2) : t \in \mathbb{R}\}$. The lower part consists of $B_s(x, r)$ with respect to $\mathbb{H}^2$ and the upper part consists of $B_s(x, r)$ with respect to $\mathbb{R}^2 \setminus \{e_2\}$. The following theorem uses this idea and gives upper bound for the radius of convexity, when the center point x is close to $\partial\mathbb{H}^2$.

Theorem 5.2. *Let $x = (x_1, x_2) \in G = \mathbb{H}^2 \setminus \{e_n\}$ with $x_2 < |x_1|$ and $r \in (0, r_0]$, where*

$$r_0 = \frac{\sqrt{x_1^2 + x_2^2} - \sqrt{2}x_2}{|x_1| + x_2}.$$

Then $B_s(x, r)$ is convex.




FIGURE 6. Metric balls $B_s(x, r)$ in $\mathbb{H}^2 \setminus \{e_2\}$ where $x = e_2/10$ and $r = 0.5, 0.75$ and 0.85 .

Proof. By simple computation we obtain that $(t^2 + 1)/2 > |t|$ and we show that $B_s(x, r_0)$ is below the curve $\{(t, |t|) : t \in \mathbb{R}\}$. Let $R = 2x_2r_0/(1 - r_0^2)$ and $y = x_1 + x_2(1 + r_0^2)/(1 - r_0^2)i$ be the Euclidean radius and center of $B_s(x, r)$. By geometry we obtain that $R = (x_1 - y_2)/\sqrt{2}$ and thus

$$\frac{2x_2r_0}{1 - r_0^2} = \frac{x_1 - x_2 \frac{1+r_0^2}{1-r_0^2}}{\sqrt{2}}$$

which implies the assertion. $\square$

We consider next the triangular ratio metric in the angular domain

$$A_\alpha = \{z \in \mathbb{R}^2 : \angle(z, 0, e_1) < \alpha/2\}, \quad \alpha \in (0, 2\pi).$$

The boundary ∂A_α consists of two half-lines, which we call sides of the domain.

Proposition 5.3. *Let $x \in A_\alpha$, $\alpha \in (0, \pi]$ and l be the line through the points 0 and $\bar{x}$. Then for all $y \in l \cap A_\alpha$ the maximal ellipse in $\overline{A_\alpha}$ touches both sides of the angular domain.*

Proof. If $\text{Im } x = 0$, then $\bar{x} = x$ and the assertion follows. We assume that $\text{Im } x < 0$. Lemma 4.3 means geometrically that the maximal ellipse in the half-plane touches the real axis on line segment between x and $\bar{y}$. Since $\angle(x, 0, e_1) = \angle(\bar{x}, 0, e_1)$ the maximal ellipse touches both sides and the assertion follows. $\square$

Lemma 5.4. *Let $G = A_\alpha$, $\alpha \in (0, \pi]$, $x \in G$ and $r \in (0, 1)$. Then*

$$B_{s_G}(x, r) = B_{s_T}(x, r) \cap B_{s_U}(x, r),$$

where T and U are the half-planes with $A_\alpha = T \cup U$. Moreover, $B_{s_G}(x, r)$ is always convex and $\partial B_{s_G}(x, r)$ is smooth for $r \leq \sin \beta / \sin(\alpha/2)$, where $\beta = \angle(x, 0, e_1)$.

Proof. By (2.2) we have $B_{s_G}(x, r) = B_{s_s}(x, r) \cap B_{s_T}(x, r)$ and by Theorem 4.4, $B_{s_G}(x, r)$ is convex as an intersection of two convex domains.

Let us denote the line through 0 and $\bar{x}$ by l . By Proposition 5.3 it is clear that $\partial B_{s_G}(x, r)$ is a circle if it does not intersect with l at two distinct points and if $\partial B_{s_G}(x, r)$ is not a circle then it is not smooth. If $\partial B_{s_G}(x, r)$ is a circle and only touches l then $\partial B_{s_G}(x, r) \cap l = \bar{x}$. We obtain that

$$s_G(x, \bar{x}) = \frac{\sin \beta}{\sin \frac{\alpha}{2}}$$

and the assertion follows. $\square$

Lemma 5.5. *Let $G = A_\alpha$, $\alpha \in (\pi, 2\pi)$ and $x \in G$. Then $B_s(x, r)$ is convex for $r \in (0, 1/2]$ and the radius $1/2$ is sharp for $\beta = \angle(x, 0, e_1) < (\alpha - \pi)/2$.*

Moreover, if $\beta > (\alpha + \pi)/2$ then $\partial B_s(x, r)$ is smooth for

$$r < \frac{\sin\left(\frac{\beta}{2} + \frac{\pi - \alpha}{4}\right)}{\sin\left(\frac{\pi + \alpha}{4} - \frac{\beta}{2}\right)}.$$

Proof. The radius of convexity $1/2$ follows from (2.2) and Theorem 3.8. Sharpness of the radius follows from the fact that if $\beta, \angle(y, 0, e_1) < (\alpha - \pi)/2$ then $s_G(x, y) = |x - y|/(|x| + |y|)$.

If $\beta > (\alpha + \pi)/2$ then by the proof of Lemma 5.4, $\partial B_s(x, r)$ is a circle if $r < \sin \beta' / \sin(\alpha'/2)$, where $\beta' = \beta/2 + (\pi - \alpha)/4$ and $\alpha' = (\pi + \alpha)/2 - \beta$. $\square$

By combining the results in the angular domain we obtain corresponding result in a polygon.

Theorem 5.6. *Let $P \subset \mathbb{R}^2$ be a polygon and $x \in P$. Then $B_s(x, r)$ is convex for all $r \in (0, 1/2]$. Moreover, if P is convex then $B_s(x, r)$ is convex for all $r \in (0, 1)$.*

Proof. The assertion follows from (2.2) and Lemmas 5.4 and 5.5. $\square$

Proof of Theorem 1.2. The assertion follows from Theorems 3.8, 4.4, 5.2 and 5.6. $\square$

Open problem 5.7. *Let $G \subsetneq \mathbb{R}^n$ be a convex domain and $x \in G$. Is $B_s(x, r)$ convex for all $r \in (0, 1)$?*

6. INCLUSION RELATIONS OF BALLS IN GENERAL DOMAINS

In this section and the following section we will consider the inclusion relations between metric balls in general domains and also some special domains.

Theorem 6.1. *Suppose that $G \subset R^n$ is a domain. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B^n \left(x, \frac{2r}{1+r} d_G(x) \right) \subset B_s(x, r) \subset B^n \left(x, \frac{2r}{1-r} d_G(x) \right).$$

Proof. We first show that $B^n(x, \frac{2r}{1+r} d_G(x)) \subset B_s(x, r)$. For each $y \in B^n(x, \frac{2r}{1+r} d_G(x))$, we have

$$\frac{|x-y|}{\min_{z \in \partial G} \{|x-z| + |y-z|\}} \leq \frac{\frac{2r}{1+r} d_G(x)}{d_G(x) + d_G(y)} \leq \frac{r}{1+r} < r$$

when $d_G(y) \geq d_G(x)$, and

$$\frac{|x-y|}{\min_{z \in \partial G} \{|x-z| + |y-z|\}} \leq \frac{\frac{2r}{1+r} d_G(x)}{d_G(x) + d_G(x) - |x-y|} \leq r$$

when $d_G(y) \leq d_G(x)$. Because y is arbitrary, we get $B(x, \frac{2r}{1+r} |x|) \subset B_s(x, r)$.

For the second inclusion $B_s(x, r) \subset B^n(x, \frac{2r}{1-r} d_G(x))$, let $y \in B_s(x, r)$ and assume that $d_G(x) < d_G(y)$. Let $z \in \partial G$ be such that $|x-z| = d_D(x)$. Then $d_G(y) \leq |y-z| \leq \frac{1+r}{1-r} |x-z| = \frac{1+r}{1-r} d_G(x)$, which implies that

$$|x-y| \leq r(|x-z| + |y-z|) \leq \frac{2r}{1-r} d_G(x).$$

□

Theorem 6.2. *Suppose that $G \subset R^n$ is a domain. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_j(x, \log(1+2r)) \subset B_s(x, r) \subset B_j \left(x, \log\left(1 + \frac{2r}{1-3r}\right) \right)$$

and the first inclusion is sharp if there exists some points w in $\partial B_s(x, r)$ such that $d_G(x) = d_G(w)$ and

$$\partial G \cap S^{n-1}(w, d_G(w)) \cap S^{n-1}(x, d_G(x)) \neq \emptyset.$$

Proof. The second inclusion follows from Theorem 6.1 and [Vu3, (3.9)]. So we only need to prove the first inclusion. For given $x \in G$ and $y \in B_j(x, \log(1+2r))$, we have $|x-y| \leq 2r \min\{d_G(x), d_G(y)\}$. Then

$$\frac{|x-y|}{\min_{z \in \partial G} \{|x-z| + |y-z|\}} \leq \frac{|x-y|}{2d_D(x)} \leq r.$$

For the sharpness: if there exists some points w in $\partial B_s(x, r)$ such that $d_G(x) = d_G(w)$ and

$$\partial G \cap S(w, d_G(w)) \cap S(x, d_G(x)) \neq \emptyset,$$

then let $z \in \partial G \cap S(w, d_G(w)) \cap S(x, d_G(x)) \neq \emptyset$. Hence

$$|x - w| \leq r(|x - z| + |w - z|) = 2rd_G(x),$$

which implies $j_G(x, w) = \log(1 + 2r)$. □

By Theorems 6.1 and 6.2 and [Vu3, (3.9)], we get the following Corollary.

Corollary 6.3. *Suppose that $G \subset R^n$ is a domain. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_k(x, \log(1 + 2r)) \subset B_s(x, r) \subset B_k\left(x, \log\left(1 + \frac{2r}{1 - 3r}\right)\right).$$




FIGURE 7. This picture is in $R^2 \setminus \{0\}$ and the boundary of the s -metric disk in Theorem 7.1 visualized as the dashed curve. Here $x = 0.4 - 0.5I$ and $r = 0.5$.

7. INCLUSION RELATIONS OF BALLS IN SOME SPECIAL DOMAINS

First we consider the punctured spaces and get what follows.

Theorem 7.1. *Let $G = R^n \setminus \{0\}$. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B\left(x, \frac{2r}{1+r}|x|\right) \subset B_s(x, r) \subset B\left(x, \frac{2r}{1-r}|x|\right).$$

Moreover, the inclusions are sharp (see Figure(7)).

Proof. The inclusions follow from Theorem 6.1. Sharpness of the first inclusion: choose $y = (1 - \frac{2r}{1+r})x$. Obviously, $|x - y| = \frac{2r}{1+r}|x|$ and $\frac{|x-y|}{|x|+|y|} = r$.

Sharpness of the second inclusion: choose $y = (1 + \frac{2r}{1-r})x$. Obviously, $|x - y| = \frac{2r}{1-r}|x|$ and $\frac{|x-y|}{|x|+|y|} = r$. □

Theorem 7.2. *Let $G = \mathbb{R}^n \setminus \{0\}$. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_j(x, m) \subset B_s(x, r) \subset B_j(x, M),$$

where $m = \log(1 + 2r)$, and $M = \log(1 + \frac{2r}{1-r})$. Moreover, the inclusions are sharp (see Figure (8)).

Proof. The first inclusion follows from Theorem 6.2.

Sharpness: Choose $y \in \partial B_s(x, r)$ with $|y| = |x|$. Then $|x - y| = 2r|x|$, whence

$$\log \left(1 + \frac{|x - y|}{|x|} \right) = \log(1 + 2r).$$

For the proof of the second inclusion, let $y \in B_s(x, r)$ with $|x| \leq |y|$.




FIGURE 8. This picture is in $\mathbb{R}^2 \setminus \{0\}$ and the boundary of the s -metric disk in Theorem 7.2 visualized as the dashed curve. Here $x = 0.5$ and $r = 0.4$.

Then

$$|x - y| \leq r(|x| + |y|)$$

and

$$|y| \leq \frac{1+r}{1-r}|x|.$$

Hence

$$\log \left(1 + \frac{|x - y|}{|x|} \right) \leq \log \left(1 + \frac{2r}{1-r} \right).$$

The case $y \in B_s(x, r)$ with $|x| \geq |y|$ follows from the fact that $s_G(x, y)$ and $j_G(x, y)$ are invariant under inversion about origin.

Sharpness: Choose $y = \frac{1+r}{1-r}x$. Then on one hand $\frac{|x-y|}{|x|+|y|} = r$, which implies $y \in \partial B_s(x, r)$. On the other hand, $\log(1 + \frac{|x-y|}{|x|}) = \log(1 + \frac{2r}{1-r})$ implies $y \in \partial B_j(x, r)$ which gives the sharpness. □

Proof of Theorem 1.3. The assertion follows from Theorems 6.1, 6.2, 7.1 and 7.2. $\square$

By Theorem 7.1 and [KV, Theorem 3.3], the following holds.

Corollary 7.3. *Let $G = \mathbb{R}^n \setminus \{0\}$. (1) For each $x \in G$ and $r \in (0, 1)$, we have $B_k(x, \log(1 + 2r)) \subset B_s(x, r)$.*

(2) For each $x \in G$ and $r \in (0, \frac{1}{2})$, we have

$$B_s(x, r) \subset B_k(x, 2 \arcsin \frac{r}{1-r}).$$

Conjecture 7.4. *Let $G = \mathbb{R}^n \setminus \{0\}$. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_k(x, m) \subset B_s(x, r) \subset B_k(x, M),$$

where $m = 2 \arcsin r$, and $M = \log(1 + \frac{2r}{1-r})$. Moreover, the inclusions are sharp (see figure (9)).




FIGURE 9. This picture is in $\mathbb{R}^2 \setminus \{0\}$ and the boundary of the s -metric disk in Conjecture 7.4 visualized as the dashed curve. In the left picture $x = 2$, $r = 0.7$, in the right one, $x = 7$ and $r = 0.7$.

Idea for the proof of Conjecture 7.4. The first part. For $y \in B_k(x, m)$, let $\alpha = \angle(x0y)$. Then $\alpha \leq \sqrt{m^2 - \log^2(\frac{|x|}{|y|})}$, and it suffices to prove

$$s_G(x, y) < \sin \frac{m}{2},$$

that is

$$\frac{|x - y|^2}{(|x| + |y|)^2} \leq \frac{|x|^2 + |y|^2 - 2|x||y| \cos \sqrt{m^2 - \log^2(\frac{|x|}{|y|})}}{|x|^2 + |y|^2 + 2|x||y|} \leq \sin^2 \frac{m}{2}$$

and thus only need to prove the following:

$$\begin{aligned} f(t) &= \left(1 - \sin^2 \frac{m}{2}\right) + \left(1 - \sin^2 \frac{m}{2}\right) t^2 \\ &\quad - 2 \left(\sin^2 \frac{m}{2} + \cos \sqrt{m^2 - \log^2 t} \right) t \leq 0, \end{aligned}$$

where $m \in [0, \pi/2]$ and $t \in [0, e^m]$.

The second part. For $y \in B_s(x, r)$, let $\alpha = \angle(x0y)$. Then

$$\begin{aligned} \alpha &= \arccos \frac{|x|^2 + |y|^2 - |x - y|^2}{2|x||y|} \\ &\leq \arccos \frac{(1 - r^2)|x|^2 + (1 - r^2)|y|^2 - 2r^2|x||y|}{2|x||y|}. \end{aligned}$$

It suffices to prove

$$\begin{aligned} f(t) &= \arccos^2 \left(\frac{1 - r^2}{2} t + \frac{1 - r^2}{2t} - r^2 \right) + \log^2 t \leq f \left(\frac{1 + r}{1 - r} \right) \\ &= f \left(\frac{1 - r}{1 + r} \right), \end{aligned}$$

where $r \in (0, 1]$ and $t \in (\frac{1-r}{1+r}, \frac{1+r}{1-r})$.




FIGURE 10. This picture is in H^2 and the boundary of the s -metric disk in Theorem 7.6 visualized as the dashed curve. Here $x = 2I$ and $r = 0.5$.

For $x \in G = H^n$, by [Vu3, (2.11)] we have $B_k(x, r) = B^n(z, |x| \sinh r)$ with $z = |x|e_n \cosh r$, and we know that $B_s(x, r) = B^n(x - e_n x_n(1 - \frac{1+r^2}{1-r^2}), \frac{2x_n r}{1-r^2})$, then it is easy to get the following Lemma.

Lemma 7.5. *Let $x \in G = H^n$ and $r \in (0, 1)$. Then*

$$B_s(x, r) = B_k(x, \log(1 + \frac{2r}{1-r})).$$

Theorem 7.6. *Let $G = H^n$. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_j(x, m) \subset B_s(x, r) \subset B_j(x, M),$$

where $m = \log(1 + \frac{2r}{\sqrt{1-r^2}})$, and $M = \log(1 + \frac{2r}{1-r})$. Moreover, the inclusions are sharp (see figure (10)).

Proof. It follows from Lemma 7.5 and [KV, Theorem 4.2] and the fact that $B_k(x, r) \subset B_j(x, r)$. □

Conjecture 7.7. *Let $G = B^n$. For each $x \in G$ and $r \in (0, 1)$, we have*

$$B_j(x, m) \subset B_s(x, r) \subset B_j(x, M),$$

where $m = \log(1 + 2r)$, and $M = \log(1 + \frac{2r}{1-r})$. Moreover, the second inclusion is sharp.

REFERENCES

- [AST] V.V. ASEEV, A.V. SYCHĚV, A. V., A.V. TETENOV: *Möbius-invariant metrics and generalized angles in Ptolemaic spaces*. (Russian) Sibirsk. Mat. Zh. 46 (2005), no. 2, 243–263; translation in Siberian Math. J. 46 (2005), no. 2, 189–204.
- [B] A. BARRLUND: *The p -relative distance is a metric*. SIAM J. Matrix Anal. Appl. **21** (1999), no. 2, 699–702 (electronic).
- [H1] P. HÄSTÖ: *A new weighted metric: the relative metric I*. J. Math. Anal. Appl. **274**, (2002), 38–58.
- [K11] R. KLÉN: *Local convexity properties of j -metric balls*. Ann. Acad. Sci. Fenn. Math. **33** (2008), 281–293.
- [K12] R. KLÉN: *Local convexity properties of quasihyperbolic balls in punctured space*. J. Math. Anal. Appl. **342** (2008) 192–201.
- [K13] R. KLÉN: *On hyperbolic type metrics*. Dissertation, University of Turku, Helsinki, 2009. Ann. Acad. Sci. Fenn. Math. Diss. No. 152 (2009), 49 pp.
- [K14] R. KLÉN: *Close-to-convexity of quasihyperbolic and j -metric balls*. Ann. Acad. Sci. Fenn. Math. **35** (2010), 493–501.
- [KLVW] R. KLÉN, H. LINDÉN, M. VUORINEN, G. WANG: *Visual angle metric and Möbius transformations*. arXiv:1208.2871 [math.MG].
- [KRT] R. KLÉN, A. RASILA, J. TALPONEN: *Quasihyperbolic geometry in Euclidean and Banach spaces*. J. Anal. **18** (2010), 261–278.
- [KV] R. KLÉN AND M. VUORINEN: *Inclusion relations of hyperbolic type metric balls*. Publ. Math. Debrecen **81** (2012), 289–311.
- [KV2] R. KLÉN AND M. VUORINEN: *Inclusion relations of hyperbolic type metric balls II*. arXiv:1105.1231 [math.MG].
- [L] H. LINDÉN: *Quasihyperbolic geodesics and uniformity in elementary domains*. Dissertation, University of Helsinki, Helsinki, 2005. Ann. Acad. Sci. Fenn. Math. Diss. No. 146 (2005), 50 pp.
- [MO] G.J. MARTIN, B.G. OSGOOD: *The quasihyperbolic metric and the associated estimates on the hyperbolic metric*. J. Anal. Math. **47** (1986), 37–53.

- [MV] O. MARTIO, J. VÄISÄLÄ: *Quasihyperbolic geodesics in convex domains II*. Pure Appl. Math. Q. 7 (2011), 379–393.
- [RT] A. RASILA, J. TALPONEN: *Convexity properties of quasihyperbolic balls on Banach spaces*. Ann. Acad. Sci. Fenn. Math. 37 (2012), 215–228.
- [Va1] J. VÄISÄLÄ: *Quasihyperbolic geometry of domains in Hilbert spaces*. Ann. Acad. Sci. Fenn. Math. 32 (2007), 559–578.
- [Va2] J. VÄISÄLÄ: *Quasihyperbolic geometry of planar domains*. Ann. Acad. Sci. Fenn. Math. 34 (2009), 447–473.
- [Vu1] M. VUORINEN: *Conformal invariants and quasiregular mappings*. J. Anal. Math. 45 (1985), 69–115.
- [Vu2] M. VUORINEN: *Metrics and quasiregular mappings*. Proc. Int. Workshop on Quasiconformal Mappings and their Applications, IIT Madras, Dec 27, 2005 - Jan 1, 2006, ed. by S. Ponnusamy, T. Sugawa and M. Vuorinen, *Quasiconformal Mappings and their Applications*, Narosa Publishing House, New Delhi, India, 291–325, 2007.
- [Vu3] M. VUORINEN: *Conformal Geometry and Quasiregular Mappings*, Lecture Notes in Mathematics 1319, Springer-Verlag, Berlin–Heidelberg–New York, 1988.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU,
FIN-20014 TURKU, FINLAND

E-mail address: `samhok@utu.fi`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU,
FIN-20014 TURKU, FINLAND

E-mail address: `ripekl@utu.fi`

DEPARTMENT OF MATHEMATICS, HUNAN NORMAL UNIVERSITY, CHANGSHA,
HUNAN 410081, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `yaxiangli@163.com`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU,
FIN-20014 TURKU, FINLAND

E-mail address: `vuorinen@utu.fi`